

Multiwavelength observations of the M 15 intermediate velocity cloud

J. V. Smoker¹, L. M. Haffner², F. P. Keenan¹, R. D. Davies³, D. Pollacco¹

¹*Astrophysics and Planetary Science Division, Department of Pure and Applied Physics, The Queen's University of Belfast, University Road, Belfast, BT7 1NN, U.K.*

²*Dept of Astronomy, University of Wisconsin, 5534 Sterling Hall, 475 North Charter Street, Madison, WI 53706, U.S.A.*

³*Jodrell Bank Observatory, University of Manchester, near Macclesfield, Cheshire, SK11 9DL, U.K.*

Accepted Received in original form

ABSTRACT

We present Westerbork Synthesis Radio Telescope H I images, Lovell telescope multibeam H I wide-field mapping, William Herschel Telescope longslit echelle Ca II observations, Wisconsin H α Mapper (WHAM) facility images, and IRAS ISSA 60 and 100 micron coadded images towards the intermediate velocity cloud (IVC) at +70 km s^{−1}, located in the general direction of the M 15 globular cluster. When combined with previously-published Arecibo data, the H I gas in the IVC is found to be clumpy, with a peak H I column density of $\sim 1.5 \times 10^{20}$ cm^{−2}, inferred volume density (assuming spherical symmetry) of ~ 24 cm^{−3} / D (kpc), and maximum brightness temperature at a resolution of $81'' \times 14''$ of ~ 14 K. The major axis of this part of the IVC lies approximately parallel with the Galactic plane, as does the low velocity H I gas and IRAS emission. The H I gas in the cloud is warm, with a minimum value of the full width half maximum (FWHM) velocity width of 5 km s^{−1} corresponding to a kinetic temperature, in the absence of turbulence, of ~ 540 K. From the H I data, there are indications of two-component velocity structure. Similarly, the Ca II spectra, of resolution 7 km s^{−1}, also show *tentative* evidence of velocity structure, perhaps indicative of cloudlets. Assuming there are no unresolved narrow-velocity components, the mean values of $\log_{10}(N(\text{Ca II K}) \text{ cm}^{-2}) \sim 12.0$ and $\text{Ca II}/\text{H I} \sim 2.5 \times 10^{-8}$ are typical of observations of high Galactic latitude clouds. This compares with a value of $\text{Ca II}/\text{H I} > 10^{-6}$ for IVC absorption towards HD 203664, a halo star of distance 3 kpc, some 3.1 degrees from the main M 15 IVC condensation. The main IVC condensation is detected by WHAM in H α with central LSR velocities of ~ 60 – 70 km s^{−1}, and intensities uncorrected for Galactic extinction of up to 1.3 Rayleigh, indicating that the gas is partially ionised. The FWHM values of the H α IVC component, at a resolution of 1 degree, exceed 30 km s^{−1}. This is some 10 km s^{−1} larger than the corresponding H I value at similar resolution, and indicates that the two components may not be mixed. However, the spatial and velocity coincidence of the H α and H I peaks in emission towards the main IVC component is qualitatively good. If the H α emission is caused solely by photoionisation, the Lyman continuum flux towards the main IVC condensation is $\sim 2.7 \times 10^6$ photons cm^{−2} s^{−1}. There is not a corresponding IVC H α detection towards the halo star HD 203664 at velocities exceeding ~ 60 km s^{−1}. Finally, both the 60 and 100 micron IRAS images show spatial coincidence, over a $0.675^\circ \times 0.625^\circ$ field, with both low and intermediate velocity H I gas (previously observed with the Arecibo telescope), indicating that the IVC may contain dust. Both the H α and tentative IRAS detections discriminate this IVC from High Velocity Clouds although the H I properties do not. When combined with the H I and optical results, these data point to a Galactic origin for at least parts of this IVC.

Key words: ISM: general – ISM: clouds – ISM: individual objects: Complex gp – ISM: structure – globular clusters: individual: M 15 – radio lines: ISM.

1 INTRODUCTION

The study of intermediate velocity clouds (IVCs) remains one of the most challenging in contemporary Galactic astronomy, with several issues concerning IVCs remaining unresolved. These include, but are not limited to, the method of their formation, their relationship (if any) with high velocity clouds (HVCs), and the question as to whether IVCs are sites of star formation in the halo of the Galaxy (Kuntz & Danly 1996; Christodoulou, Tohline & Keenan 1997; Ivezić & Christodoulou 1997). This latter possibility is underpinned by the fact that within the Galactic halo, there exists a population of early B-type stars whose velocities, ages and distances from the Galactic plane (z) are incompatible with them being formed within the disc. A possible site for their formation is IVCs/HVCs via cloud-cloud collisions and subsequent compression of the gas (Dyson & Hartquist 1983). Such collisions are thought to be a viable star formation mechanism within at least the discs of galaxies, albeit where the gas density and cloud-cloud collision rates are somewhat higher than inferred in IVCs/HVCs (Tan 2000).

The solution to both the star formation question and also any possible relationship between HVCs and IVCs requires both the analysis of aggregate parameters of well-defined samples of IVCs and HVCs, and also more detailed studies of individual objects. In this paper we report on radio H I aperture synthesis, H I multibeam wide field mapping, longslit Ca II observations, Wisconsin H α Mapper (WHAM) facility images, and IRAS sky-survey archive data retrieval towards a particular IVC located in the general direction of the M 15 globular cluster (RA=21^h 29^m 58.29^s, Dec=+12° 10′ 00.5″ (J2000); $l=65.01^\circ$, $b=-27.31^\circ$). These observations are amongst the first H I synthesis data to be taken of positive-velocity IVCs, which remain poorly-studied as a group of objects.

The M 15 H I cloud lies at a velocity of $\sim +70$ km s⁻¹ in the dynamical Local Standard of Rest (Allen 1973); its distance tentatively lies between ~ 0.8 –3 kpc (Little et al. 1994; Smoker et al. 2001a). The upper distance limit is gleaned from the fact that IVC absorption at $\sim +70$ to $+80$ km s⁻¹ is observed in the spectrum of HD 203664, a halo star of distance 3 kpc and ~ 3.1 degrees from M 15 (Little et al. 1994), combined with the detection of IV H I approximately mid-way between the M 15 IVC and HD 203664.

The deviation velocity of the M 15 IVC at such a mid-Galactic latitude puts it in on the borderline between the normal definitions for intermediate and high velocity clouds (c.f. Fig. 1 of Wakker 1991), although in common with Sembach (1995) and Kennedy et al. (1998) here we classify it as an IVC. The line-of-sight position of the M 15 IVC is between the negative-velocity Local Group barycentre cloud Complex G and the Galactic centre clouds (Fig. 8 of Blitz et al. 1999), hence the M 15 IVC is a part of IVC Complex gp (Wakker 2001).

Previous observations in H I emission using the Lovell and Arecibo telescopes (Kennedy et al. 1998; Smoker et al. 2001a) have shown that the IVC consists of several condensations of gas spread out over an area of more than 3 square degrees, with structure existing down to the previous resolution limit of ~ 3 arcmin. The brightest component is located towards M 15 itself and has peak H I column density at the Arecibo resolution of $\sim 8 \times 10^{19}$ cm⁻². In this paper,

we study this part of the IVC at higher spatial resolution. The mass of this particular clump is $\sim 20 D^2 M_\odot$, (where D is the distance in kpc), thus for this particular object, in the absence of H₂, there is insufficient neutral gas to form an early-type star. Low-resolution absorption-line Ca II and Na I spectroscopy (Lehner et al. 1999) towards cluster stars tentatively found cloud structure (or variations in the relative abundance) over scales as small as a few arcsec, with fibre-optic array mapping in the Na I D absorption lines (Meyer & Lauroesch 1999) obtaining similar results with structure visible on scales of ~ 4 arcsec (velocity resolution ~ 16 km s⁻¹). Using empirical relationships between the sodium and hydrogen column densities, Meyer & Lauroesch (1999) derived values of the H I column density towards the cluster centre of $\approx 5 \times 10^{20}$ cm⁻², some 6 times higher than that found using the Arecibo H I data alone; the difference may be attributable to fine-scale cloud structure. Assuming spherical symmetry, a volume density of ≈ 1000 cm⁻³ is implied by these latter results, similar to values obtained for gas in the local ISM (e.g. Faison et al. 1998, although see Lauroesch, Meyer & Blades 2000). Such a high volume density and implied overpressure with respect to the ISM perhaps indicates that the assumption of spherical symmetry is invalid and that there may be some sheet like geometry in the IVC as has been postulated for low-velocity gas (Heiles 1997).

In the current paper, we extend our studies of the M 15 IVC to higher resolution and different wavelength regions in order to investigate three areas. Firstly, H I synthesis mapping, WHAM H α and IRAS ISSA survey data retrieval towards the IVC were performed to see if the H I, H α and infrared properties of the M 15 IVC are compatible with either low velocity gas or HVCs in general, and whether there are any differences between the types of object, perhaps attributable to differences in the formation mechanism or current environment (for example, distance from the ionising field of the Galaxy). Secondly, wide-field medium-resolution H I data were obtained in search of more IVC components, to trace how the gas kinetic temperature changes with sky position, and possibly determine the relative distance of cloud components from the Galactic plane (c.f. Lehner 2000). Thirdly, longslit echelle Ca II observations were undertaken, using the centre of M 15 as a background continuum source, in order to look for small-scale velocity and column density substructure within the IVC which could indicate the presence of cloudlets, collisions between which in certain IVCs may be responsible for star formation in the Galactic halo.

Section 2 describes the observations and data reduction, Sect. 3 gives the results, Sect. 4 contains the discussion and Sect. 5 presents a summary and the conclusions.

2 OBSERVATIONS AND DATA REDUCTION

2.1 Westerbork aperture synthesis H I observations

21-cm aperture synthesis H I observations of the M 15 IVC were obtained during two observing sessions, each of 12 hours, using the Westerbork Synthesis Radio Telescope (WSRT). The first in 19 December 1998 had a minimum antenna spacing of 32-m, the second in 17 April 1999 had a corresponding separation of 72-m. The velocity resolution for all observations was 1.03 km s⁻¹.

Standard methods within AIPS^{*} were used to reduce the visibility data. Reduction included amplitude calibration using 3C 286 and 3C 48 (assuming flux densities of 14.76 Jy and 15.98 Jy, respectively), phase calibration and flagging of bad data, and concatenation of the two UV datasets using DBCON. The calibrated dataset was mapped with IMAGR using both quasi-natural and uniform weightings, with the Arecibo map from Smoker et al. (2001a) being used to set the locations of the CLEAN boxes for each velocity channel interactively. The respective beamsizes of the final images were $111'' \times 56''$ and $81'' \times 14''$ (approximately North–South by East–West), with the corresponding rms noises being 2.3 mJy beam⁻¹ and 1.3 mJy beam⁻¹ for the naturally and uniformly-weighted data.

As the WSRT maps suffer from missing short-spacing information, it was decided to create ‘total power’ channel maps by combining the current observations with the single dish Arecibo data. For this we used MIRIAD (Sault, Teuben & Wright 1995). The two datasets were first regridded so that they had the same central coordinate and channel width. Following this, we multiplied the Arecibo data by the WSRT single-dish beam, converted to flux density units, and combined the WSRT and weighted Arecibo maps using IMMERGE. Combination was performed by specifying an annulus of between 50–150-m in the Fourier domain where the high and low resolution images were made to agree. Within this annulus it was found that, where the H I was strong, the flux density scales of the two datasets were the same. Hence the ‘calibration factor’ used when combining the two datasets was set to unity. The resulting combined channel maps were divided by the WSRT single dish beam to produce the final combined images. These are of the resolution of the WSRT data ($111'' \times 56''$), but with the correct zero-spacing flux. Finally, moments analysis and Gaussian fitting were performed on the WSRT channel maps and combined cube from +50 to +90 km s⁻¹ in order to determine the H I column densities, velocities and velocity widths of the IVC. H I column densities were derived using the relationship $N_{\text{HI}} = 1.823 \times 10^{18} \int T_{\text{B}} dv$ (cm⁻²), where v is the velocity in km s⁻¹.

2.2 Lovell Telescope multibeam H I observations

Lovell telescope multibeam 21-cm H I observations covering a $\sim 4.9 \times 9.3$ degree RA–Dec region, centred upon RA=21^h22^m47^s, Dec=+10°00′29″ (J2000) were taken during 2 June 2000. The resolution of these data is 12 arcmin spatially and 3.0 km s⁻¹ in velocity. The total integration time was 4 hours and the RMS noise per channel was ~ 0.1 K. The observed field is not quite fully sampled, with the spacing between the beams being 10 arcmin, compared with the spatial resolution of 12 arcmin. During observing, data were calibrated in terms of antenna temperature by a frequently-fired noise diode. Off-line, bootstrapping of the flux scale to some previous Lovell telescope H I observations taken during 1996 enabled calibration in terms of brightness temperature (T_{B}). Data reduction procedures will be




Figure 1. Lovell telescope multibeam H I spectrum towards RA=21^h26^m47^s, Dec=+09°16′23″ (J2000). The dotted line shows the results of a single-component Gaussian fit. The brightness temperature calibration is accurate to ~ 40 per cent with the negative features present near to ~ 0 km s⁻¹ being an artefact of the baseline removal procedure. The dashed line is the (data–single Gaussian fit) residual.

discussed in more detail in a future paper. For now, we simply note that most of the process is automated, the bandpass removal being performed within AIPS⁺⁺ using LIVEDATA and the data being gridded using GRIDZILLA (Barnes 2001). As the bandpass is simply a running average of the observed spectra in a window, incorrect removal is performed around the low-velocity gas where there is no ‘empty’ region of the sky. However, at the velocity of the IVC ($v \sim +70$ km s⁻¹) the baseline subtraction is adequate. An example of a spectrum after automated bandpass removal is shown in Fig. 1.

After gridding, the data were imported into AIPS whence a column density map was made of the entire field between +50 and +90 km s⁻¹ which is presented in section 3.2.

2.3 William Herschel Telescope longslit UES observations

WHT Utrecht Echelle Spectrograph (UES) observations towards the centre of the M15 globular cluster were taken during bright time on 19–21 July 2000. Twin EEV CCDs were used. In combination with the E79 grating, this gave near-complete wavelength coverage from 3800–6300 Å, with a slitwidth of 0.85″ providing an instrumental FWHM resolution of ~ 6 km s⁻¹. On the first night of the run, a slit height of 16 arcseconds was used; this has the advantage that the full wavelength coverage is obtained with no overlapping orders on either of the CCDs. On the second and third night’s observations, in order to maximise the amount of sky available for the calcium lines at 3933.66 Å, a slit of height 22 arcseconds was used which leads to overlapping orders in the red CCD. ThAr arcs were taken two or three times a night to act as the wavelength calibration with Tungsten and sky flats also being obtained. The echelle proved

^{*} AIPS is distributed by the National Radio Astronomy Observatory, U.S.A.

to be remarkably stable with the centres of the arc lines varying by less than 1.5 pixels throughout the whole of the run.

The original intention of the observations was to obtain 7 parallel cuts through the IVC, separated by 0.7 arcsec, and use these to make up an RA-Dec rectangle of size $\sim 5'' \times 20''$ within which cloudlet sizes could be obtained. Unfortunately, due to the presence of Sahara dust and non-ideal seeing conditions (1.5''–2.0''), integration times were somewhat longer than had been hoped for and in the end only two longslit positions were obtained, offset by a position angle of 85 degrees. Additionally, on the second night we obtained 1200-s of integration towards a blank sky region with a 0.85'' slit and on the third night a total of 5×400-s, interleaved with the object exposures and using a slit of width 4.0'', corresponding to ~ 9500 -s of integration with a slit of 0.85'' used for the IVC observations.

Data reduction was performed using standard methods within IRAF[†]. The aims were firstly to obtain a spectrum of the inner part of the globular cluster over the whole wavelength region, and more importantly, produce longslit spectra of the Ca II lines alone for the two longslit positions taken. Data reduction included debiasing, and use of the sky flatfields to get rid of the vignetting that was clearly visible on many of the orders. Cosmic ray hits near the Ca II line were removed within FIGARO (Shortridge et al. 1999) using CLEAN. Because of the small pixel-to-pixel variation and pixel size ($\sim 0.2''$), no flatfielding to remove the pixel-to-pixel variation was performed on the data. The images were rotated in order to make the cross-dispersion axis occur along the image rows, after which sky subtraction was performed. As expected, this proved to be challenging.

2.3.1 Sky subtraction

Two methods were used to estimate the sky value and the results compared. The first used the data from the third nights observation where the blank sky exposures were interleaved with the object exposures. For these data, we scaled the high signal-to-noise combined sky flat (taken at twilight with the 0.85 arcsec slit) to produce an image ‘scaled_sky’ and then smoothed this in wavelength to a resolution equivalent to a 4 arcsecond slit. The scaling factor was chosen so that the resulting image was the same as the blank sky image taken with the 4 arcsecond slit for an equivalent of 5400-s of integration. The image ‘scaled_sky’ was then subtracted from each 5400-s of the object data. This method has the advantage that in the blank sky images there are clearly no contaminating counts from the object, but is susceptible to changes in the observing conditions and requires careful removal of the bias strip in both sky and object frames. Fig. 2 shows a cut across the dispersion axis for the sky exposures on night 3. As can be seen, there are few sky counts, although obviously in the outer parts of the slit there are few object counts too, so errors in the equivalent width increase rapidly in the extracted spectra for this region. For an equivalent 5400-s of integration time, at $\lambda \sim 3937\text{\AA}$, there were ~ 3 ADU sky counts and somewhat less at $\lambda=3933.66\text{\AA}$,




Figure 2. Cut across the dispersion axis at $\sim 3937\text{\AA}$ of a blank sky region, equivalent to 5400-s of object observing time on night 3. The ‘dips’ in the spectrum correspond to the interorder gaps.

due to the ‘sky’ effectively being a solar (G-type) spectrum which has strong Ca II absorption lines.

The method of sky subtraction that we had envisaged performing *a priori* involved removal of sky using parts of the longslit that appeared to be free of emission from the globular cluster. The problem with this method is illustrated by Fig. 3(a), which shows cuts across the dispersion axis for 5400-s of data taken on the third night at each of the two slit positions. As is apparent from the Figures, the ‘sky’ level at the same position on the order at the two different slit positions is different by ~ 3 –4 ADU per 5400-s of integration. During night 3, at $\lambda \sim 3937\text{\AA}$, position p1 had ~ 8 ADU as its ‘sky’ value, with the corresponding value for p2 being ~ 12 ADU. These clearly are much larger than the values obtained by exposing on the night sky and combined with the fact that p1 and p2 are different indicates that we are still obtaining counts from the cluster in the outer parts of the slit. This is confirmed by performing an extraction using such counts; in some parts of the spectrum the IVC absorption becomes negative which is unphysical.

Because of the lack of object-free emission on the slit, we decided to remove the sky by subtracting the smoothed version of the sky images from our object data. This method relies upon the sky conditions being similar throughout the three nights of observing. This was checked by taking cuts across three places on the dispersion axis of each 5400-s worth of data and checking for variability. At $\lambda \sim 3937\text{\AA}$, the values varied by only 2 ADU per 5400-s between nights 1 and 3 for the two slit positions. This variability, combined with the error in removing the bias, gives us an error in the final sky value of ~ 4 ADU at $3937\text{\AA}$, compared with a *peak* continuum value at this point of ~ 30 ADU.

2.3.2 Extraction and Analysis

The sky-subtracted spectra were extracted within FIGARO by summing over each 10 columns, corresponding to the worst seeing of ~ 2.0 arcseconds. Wavelength calibration was then undertaken, taking into account the shift in the λ scale along the length of the slit. Finally, normalisation and profile fitting were performed within DIPSO (Howarth et al. 1996). The final spectra have spatial resolution of ~ 2 arcseconds and velocity resolution of $\sim 7\text{ km s}^{-1}$.

[†] IRAF is distributed by the National Optical Astronomy Observatories, U.S.A.




Figure 3. (a) Cut across the dispersion axis at $\sim 3937\text{\AA}$ for the two slit positions with 5400-s of integration, showing the difficulty in determining the sky. IOG refers to the interorder gap between the echelle orders (b) Cut along the dispersion axis before normalisation.

Data were analysed using standard profile fitting methods within DIPSO using the ELF and IS programs. Before fitting, the stellar line was removed by fitting the profile by eye to the lower-wavelength part of the spectrum up to the peak stellar absorption unaffected by LV and IV gas, and creating a mirror of this spectrum for the higher wavelength data. The derived stellar spectrum has a stark-like line profile, typical of stellar Ca II lines, and was subtracted from the whole spectrum to leave the interstellar components only. After stellar-line removal, the ELF routines were used to fit Gaussian profiles to input spectra and hence provide the equivalent width, peak absorption and full width half maximum velocity values for the low velocity gas and IVC components. These results were compared to the theoretical absorption profiles computed by the IS suite of programs within DIPSO. These theoretical profiles are derived from the observed b -values (corrected for instrumental broadening), atomic data for the Ca II lines taken from Morton (1991) and initial guesses for the Ca II column density. Input parameters were varied until a good fit was produced to match the profiles obtained using ELF. This method is only appropriate in the regime where the lines are not saturated and are resolved in velocity; both caveats apply for a number of positions in the current dataset, although the presence of narrow, unresolved components cannot be ruled out. If present, these would cause our estimated column densities to be too low. The final ELF and IS fits provide the Ca II number densities and values for the FWHM velocities at each of the positions along the slit.

2.4 Wisconsin H α Mapper observations

Data were retrieved from the Wisconsin H α Mapper (WHAM) survey in the range $+06^\circ < \text{Dec.} < +15^\circ$ and $20^h20^m < \text{RA} < 21^h38^m$ (J2000). These data are of resolution 1 degree spatially and $\sim 12 \text{ km s}^{-1}$ in velocity, with a velocity coverage at this sky position from ~ -120 to




Figure 4. WSRT H I column density map of the M15 IVC integrated between $+50$ and $+90 \text{ km s}^{-1}$ and at a resolution of $111'' \times 56''$ ($\sim$ NS by EW). Contour levels are at $N_{\text{HI}} = (2, 4, 6, 8, 10) \times 10^{19} \text{ cm}^{-2}$.

$+90 \text{ km s}^{-1}$ in the LSR. Data were reduced using standard methods, which included conversion into the LSR and removal of the geocoronal H α line that appears at velocities in the range $\sim +30$ to $+40 \text{ km s}^{-1}$ (Haffner 1999; Haffner et al. 2001b). Due to the removal of this line, the noise in this part of the spectrum is slightly enhanced compared with the typical RMS noise value (measured for the current spectra) of $\sim 1\text{--}3 \text{ mR (km s}^{-1})^{-1}$. These values are typical for the WHAM survey as a whole. Finally, we note if the IVC H α flux originates from above the Galactic plane, then a correction for Galactic extinction of 1.25 would be necessary, calculated from a $E(B - V) = 0.10$ observed towards M15 by Durrell & Harris (1993), combined with the extinction law of Cardelli, Clayton & Mathis (1989). We have not applied this correction to our data so the IVC H α fluxes may be ~ 25 per cent too low.

2.5 IRAS ISSA data retrieval

Both the 60 and 100 micron images of a field of size $0.675^\circ \times 0.625^\circ$ were extracted from the on-line versions of the IRAS Sky Survey Atlas (ISSA). These images are of resolution ~ 5 arcmin and have most of their zodiacal emission removed although they have an arbitrary zero-point.

3 RESULTS

3.1 WSRT aperture synthesis H I results

Fig. 4 shows the H I column density map of the IVC for the low-resolution WSRT data alone, with Fig. 5 showing the corresponding map of the combined (WSRT plus Arecibo) image overlaid on the digitised Palomar Observatory Sky Atlas (POSS-I) red image regridded to a 4 arcsec pixel size. The major axis of this part of the IVC lies parallel with




Figure 5. WSRT plus Arecibo combined H I column density map of the M15 IVC integrated between $+50$ and $+90$ km s^{-1} and at a resolution of $111'' \times 56''$, overlaid on the red POSS-1 digital sky survey image towards M15. Velocities are in the LSR with contour levels being at $N_{\text{HI}} = (2, 4, 6, 8, 10, 12, 14) \times 10^{19} \text{ cm}^{-2}$. Lines of constant Galactic longitude and latitude are also plotted on the figure.

the Galactic plane. The H I channel maps of the combined WSRT plus Arecibo dataset in brightness temperature (T_B) are shown in Fig. 6, where flux density per beam is related to T_B by $S_{\text{mJy/beam}} = 0.65 \Omega_{\text{as}} T_B / \lambda_{\text{cm}}^2$; here Ω_{as} is the beam area in arcsec^2 and λ_{cm}^2 is the observed wavelength in cm (Braun & Burton 2000). Immediately obvious from each of these figures is the fact that the H I is clumpy in nature, as is seen in other intermediate and high velocity clouds observed at a similar resolution (e.g. Weiß et al. 1999; Braun & Burton 2000). Variations in the column density of a factor of ~ 4 on scales of ~ 5 arcmin are observed, corresponding to scales of $\sim 1.5 D$ pc, where D is the the IVC distance in kpc. The peak brightness temperature in the combined map (of resolution $111'' \times 56''$), is ~ 8 K, rising to ~ 14 K for the highest resolution WSRT map with beamsize $81'' \times 14''$. These values compare with peak values of T_B for HVCs of ~ 25 K towards Complex A at 2 arcmin resolution (Schwarz, Sullivan & Hulsbosch 1976), ~ 34 K observed towards Complex M at a resolution of 1 arcmin (Wakker & Schwarz 1991), and ~ 75 K observed to the compact high velocity cloud CHVC 125+41–207 (Braun & Burton 2000). Similarly the IVC observed by Weiß et al. (1999) has peak T_B exceeding 7 K, at a resolution of 1 arcmin.

The peak H I column density in the combined image is $\sim 1.5 \times 10^{20} \text{ cm}^{-2}$. If we assume that the cloud is spherically symmetric or filamentary (Stoppelenburg, Schwarz & van Woerden 1998), and using a cloud size full width half maximum (c.f. Wakker & Schwarz 1991) of ~ 7 arcmin or 2 Dpc for the brightest point, we obtain a peak volume density of $\sim 24 D^{-1} \text{ cm}^{-3}$. At the assumed distance, this value is a lower limit as it is likely that there will be more structure

on smaller scales, as indicated by the spectra of Meyer & Lauroesch (1999).

Fig. 7 shows the brightness temperature (T_B) profile at the position of peak temperature in the uniformly-weighted data, with Fig. 8 depicting the combined temperature profiles at a number of positions in the cloud. The full width half maximum (FWHM) varies from 5 km s^{-1} at the position of the three peaks in brightness temperature (Fig. 7 and Fig. 8(a)–(c)), to more than 12 km s^{-1} at other locations in the IVC (Fig. 8(d)). Some parts of the cloud (cf Fig. 8(b)) are well-fitted by a Gaussian, whereas others, such as Figs. 8(e), appear to be made up of two narrowish components. Although the signal-to-noise in this latter part of the cloud is low, both the WSRT and Arecibo datasets show evidence for a two-component velocity substructure, perhaps indicative of H I cloudlets or of overlapping clouds in the line of sight.

Fig. 9 displays the results of single-component Gaussian fitting to the IVC using XGAUSS and shows how the smaller values of FWHM (and hence implied kinetic temperature) tend to occur where the H I is brightest in the three cores. At these positions, in the absence of turbulence, the kinematic temperature is of the order of 500 K. The M15 IVC does not show any regular velocity gradients across its field as has been seen in many other HVCs and IVCs (e.g. Weiß et al. 1999; Brüns et al. 2000).

We finally note that an interesting feature of the H I surface density map is that the IVC is approximately centred upon the globular cluster, with the regions of lowest column density being located towards the globular cluster centre. It is tempting to suggest some mechanism whereby the IVC and globular cluster are associated, for example by capture of the IVC by M15, with the gas in the centre being ionised by the cluster itself. However, the existence of other IVC components in the nearby field, combined with the fact that M15 is located at a radial velocity of $\sim -100 \text{ km s}^{-1}$ (Harris 1996), compared with the the IVC at $\sim +70 \text{ km s}^{-1}$, makes it likely that the two objects are simply line-of-sight companions.

3.2 Lovell telescope multibeam H I results

The H I column density map of a 4.9×9.3 degree field, with spatial resolution 12 arcmin, centred to the South-West of M15, integrated from $+50$ to $+90 \text{ km s}^{-1}$, is shown in Fig. 10.

No new firm H I detections were obtained that are not already given in the Leiden/Dwingeloo survey (Hartmann & Burton 1997), Kennedy et al. (1998) or Smoker et al. (2001a). However, the extent of an IVC component around $\text{RA}=21^{\text{h}}27^{\text{m}}$, $\text{Dec}=+09^{\circ}10'$ (J2000; feature ‘D’) was determined to be ~ 0.7 degrees in RA–Dec at full width half power in the N_{HI} map. An example H I spectrum in this area is shown in Fig. 1. At this position, a single-component Gaussian fit within DIPSO gives a velocity centroid of $68 \pm 0.5 \text{ km s}^{-1}$, FWHM of $12.0 \pm 1.2 \text{ km s}^{-1}$ and peak T_B of ~ 1 K, similar properties to the main complex studied in this paper. The peak IVC H I column density towards this component (‘D’) is $\sim 3 \times 10^{19} \text{ cm}^{-2}$.

Also plotted on Fig. 10 are ‘A’, ‘AR’ and ‘HD’. ‘A’ corresponds to the main clump observed by Kennedy et al. (1998) which has FWHM ~ 12 – 15 km s^{-1} at 12 arcmin resolution.




Figure 6. WSRT plus Arecibo combined H I brightness temperature channel maps towards the M15 IVC. Velocities are in the LSR and the velocity resolution is 1.03 km s^{-1} . Contour levels are at $T_B = (-1, 1, 2, 4, 8)$ Kelvin. The spatial resolution is $111'' \times 54''$ ($\sim$ NS by EW).

The peak IVC column H I density derived using the current observations towards component ‘A’ of $3.7 \times 10^{19} \text{ cm}^{-2}$ is close to the value of $3.9 \times 10^{19} \text{ cm}^{-2}$ observed by Kennedy et al. (1998) with the same telescope and gives us faith in our calibration. ‘AR’ refers to previous IVC detections using the Arecibo telescope which have $\text{FWHM} = 12 \text{ km s}^{-1}$ and $v_{\text{LSR}} = +61 \text{ km s}^{-1}$ at a resolution of 3 arcmin. ‘HD’ is the position of the halo star HD 203664 (of distance 3 kpc) in which two strong interstellar IVC Ca II K absorption components are seen with $\text{FWHM} = 2.8$ and 3.2 km s^{-1} and $v \sim +80$ and $+75 \text{ km s}^{-1}$ respectively (Ryans, Sembach

& Keenan 1996). As this is an absorption-line measurement towards a star, the resolution is sub-arcsecond. Finally, we note that there is a hint of emission at $\text{RA} = 21^{\text{h}}23^{\text{m}}04^{\text{s}}$, $\text{Dec} = +14^{\circ}12'42''$ (feature ‘E’ on Fig. 10) although this is very close to the noise and may be a spurious detection.

3.3 William Herschel Telescope longslit UES results

Equivalent widths of some strong stellar lines from the inner 3 arcseconds of the first slit position are shown in Table 1.




Figure 7. WSRT uniformly-weighted H I brightness temperature profile of the M15 IVC at the position of peak T_B . The beamsize is $81'' \times 14''$. The dotted line depicts the Gaussian fit to the spectrum and has $\text{FWHM} = 5.2 \pm 0.2 \text{ km s}^{-1}$. The dashed line shows the (data – single Gaussian fit) residual.

Table 1. Equivalent widths of some strong lines towards the centre of M15. Where there are two values present, these refer to results taken on nights 1 and 2 respectively. Stellar types corresponding to the EWs for that particular line are taken from Jaschek & Jaschek (1987) (JJ87).

Species	λ_{rest}	EW (Å)	JJ87 Type
Ca II K	3933.66	1.6, 1.5	A2
Fe I	4045.82	0.16, 0.17	A3
Ca I	4226	0.25, 0.28	F2

As the core of the cluster is unresolved from the ground, the spectrum obtained is composite. Fig. 11 shows the Ca II K spectra at this position. The equivalent widths of the LV and IV interstellar components at this position are $\sim 0.3 \text{ Å}$ and $\sim 0.08 \text{ Å}$ respectively. The strength of the LV component multiplied by $\sin(b)$ of $\sim 135 \text{ mÅ}$ compares with the canonical value of $\sim 110 \text{ mÅ}$ integrated EW of the Ca II K perpendicular to the Galactic plane (Bowen 1991).

We now turn to the interstellar Ca II observations. Fig. 12 shows the extracted spectra at three and five positions along the dispersion axis for slit positions 1 and 2 respectively, where the signal-to-noise in the continuum exceeds ~ 10 . The positions shown are separated by 2 arcseconds which corresponds to the worst seeing during the run, and is also the spatial resolution to which the data were smoothed when the extraction was performed.

The prime aim of these longslit observations was to determine whether there is velocity substructure within the IVC. As can be seen from a few of the sightlines, there is *tentative* evidence for such structure, with two cloudlets being present at LSR velocities of $\sim +52 \text{ km s}^{-1}$ and $\sim +66 \text{ km s}^{-1}$ (Fig. 13). This corresponds to a separation of ~ 8 pixels so is not an artefact caused by shifts in the echelle over periods of several hours. However, the signal-to-noise of the data is low, and follow-up observations are required to con-




Figure 8. WSRT plus Arecibo combined H I brightness temperature profiles towards selected regions of the M15 IVC. (a) First T_B peak ($\text{RA} = 21^{\text{h}} 29^{\text{m}} 44^{\text{s}}$, $\text{Dec.} = +12^{\circ} 09' 42''$). (b) Second T_B peak ($\text{RA} = 21^{\text{h}} 29^{\text{m}} 36^{\text{s}}$, $\text{Dec.} = +12^{\circ} 06' 22''$). (c) Third T_B peak ($\text{RA} = 21^{\text{h}} 30^{\text{m}} 13^{\text{s}}$, $\text{Dec.} = +12^{\circ} 10' 42''$). (d) High FWHM region ($\text{RA} = 21^{\text{h}} 29^{\text{m}} 21^{\text{s}}$, $\text{Dec.} = +12^{\circ} 04' 22''$). (e) Two component profile ($\text{RA} = 21^{\text{h}} 30^{\text{m}} 38^{\text{s}}$, $\text{Dec.} = +12^{\circ} 19' 22''$). The equinox is J2000.

firm this finding. We note that although the observations of Meyer & Lauroesch (1999) may have also been expected to find such velocity substructure at their resolution of 14 km s^{-1} , their observations were in the Na I D lines, which in some cases do not show velocity components, even when the Ca II lines do (Ryans et al. 1996).

The current observations could also be used to estimate the IVC Ca II column densities along the slit as explained in section 2.3.2. Figure 14 displays the equivalent widths and Ca II column densities for a number of the slit positions for the second slit orientation. Our Ca II K equivalent widths range from 0.06 – 0.09 Å . These are slightly lower than those measured by Lehner et al. (1999) towards other parts of the IVC with low-resolution data, which are in the range 0.05 – 0.20 Å , with a median value of 0.10 Å . Towards the nearby halo star HD 203664 (which has an IVC H I column density of less than $\sim 10^{18} \text{ cm}^{-2}$), Ryans et al. (1996) measured an IVC Ca II equivalent width of $0.06 \pm 0.01 \text{ Å}$. The Ca II/H I ratio for the current data towards the M15 IVC varies from 2.1×10^{-8} to 2.9×10^{-8} . Unfortunately, poor quality of our measurements caused by low signal-to-noise,




Figure 9. WSRT plus Arecibo H I combined image. Greyscale; full width half maxima velocity widths obtained using single-component Gaussian fitting. Contours; peak values of brightness temperature of the same single-component fits with levels at (2,4,6,8) Kelvin.

and uncertainties in determining the sky and continuum levels, results in our errors being larger than the differences between these values.

3.4 Wisconsin H α Mapper (WHAM) Results

3.4.1 The main IVC towards M15

Of the 184 spectra extracted from the WHAM survey, a clearly-defined H α component at intermediate velocities exceeding $+50 \text{ km s}^{-1}$ is only obvious at three positions (A0, A2, A5) with a weak detection towards (A7). We note that because M15 itself is at an LSR velocity of $\sim -100 \text{ km s}^{-1}$, it does not contaminate the observed spectrum. The spectra observed at these positions are shown in Fig. 15(c). Fig. 15(a) depicts the locations of these WHAM pointings relative to the Lovell telescope H I surface density map, with integration of velocities from $\sim +50$ – 90 km s^{-1} in the WHAM H α data also being superimposed. Fig. 15(a) also shows the WHAM data integrated from $\sim +50$ – 90 km s^{-1} , over the whole field mapped by the multibeam observations. The gas with brightness $\sim 0.1 \text{ R}$ is quite extended about the main IVC condensation, with an additional weak signal detected towards the tentative H I detection ‘E’ (see Fig. 10).

Results of three-component Gaussian fitting at the four positions are shown in Table 2, uncorrected for Galactic extinction. These fits take into account the instrumental profile of the instrument, which is comprised of a 12 km s^{-1} Gaussian with low-level wings superimposed. Although almost equally-well fitted using just two components (for the LV and IV gas), we chose to fit using three components due to the slight asymmetry in the LV component, most clearly seen towards position (A7). Table 2 also shows the




Figure 10. Lovell telescope multibeam H I column density map of resolution 12 arcmin in the general direction of the M15 globular cluster integrated from $+50$ to $+90 \text{ km s}^{-1}$ in the LSR. Contour levels are at $(1,2,3) \times 10^{19} \text{ cm}^{-2}$, with greyscale levels from 0 – $4 \times 10^{19} \text{ cm}^{-2}$. For the meaning of the labels see Sect. 3.2.




Figure 11. Ca II K central $3''$ towards M15.




Figure 12. Normalised spectra each 2 arcseconds along the slit. Spectra are offset in units of 1.0 for clarity. (a) Ca II K spectra at slit position 1. (b) Ca II K spectra at slit position 2.




Figure 13. Solid line: Ca II K LV and IV spectrum. Dotted line; ELF multicomponent fit. Dashed line; (data-model) residual plus 1.0. A possible second intermediate velocity component is marked by IV?.




Figure 14. Ca II fit results along the second slit position.

central velocity and velocity widths observed in H I towards the component (A5), obtained by smoothing the Lovell telescope multibeam data to a resolution of 1 degree.

The spectrum with the strongest IV H α emission (A5) is depicted in Fig. 16. This is the nearest grid position to the M 15 IVC, whose centre lies approximately at $l=65.01^\circ$, $b=-27.31^\circ$. The peak IV H α brightness in this direction obtained using three-component Gaussian fitting is 0.035 Rayleighs $(\text{km s}^{-1})^{-1}$ for the low velocity gas, and 0.033 Rayleighs $(\text{km s}^{-1})^{-1}$ for the intermediate velocity gas, where 1 Rayleigh is $10^6/4\pi$ photons $\text{cm}^{-2} \text{s}^{-1} \text{sr}^{-1}$. The integrated fluxes are 2.2 ± 0.1 and 1.3 ± 0.1 Rayleighs for the (total) LV and IV gas respectively. The Gaussian fit gives centroids of -53.9 ± 4.6 , -4.4 ± 0.7 and $+64.3 \pm 0.4$ km s^{-1} , and FWHM velocity widths of 26.7 ± 10.4 , 47.8 ± 2.6 and 31.5 ± 1.4 km s^{-1} , for the low and intermediate velocity gas respectively. The velocity centroid at this position (A5) agrees within the errors with the H I data smoothed to the WHAM spatial resolution; this contrasts with the results of Tufte, Reynolds & Haffner (1998) who tentatively found an offset in velocity between H α and H I velocities of ~ 10 km s^{-1} towards various HVC complexes. For the position (A5), the velocity width of the IV H α spectrum of ~ 32 km s^{-1} is some 10 km s^{-1} greater than the H I data at 1 degree resolution. For a gas at $\sim 10^4$ K, some ~ 22 km s^{-1} of this is caused by thermal broadening, with the remaining width being due to non-thermal motions and beamsmeared of different components. The difference between the H I and H α widths may imply that the two phases are not mixed. However, at least qualitatively, there is reasonable coincidence between the H α and H I peaks (Fig. 15(a)), although the mapped area is small.

The second positive detection (towards (A2) in Fig. 15), occurs in a region where the local H I column density, at the Lovell telescope resolution of ~ 12 arcmin, is lower than $1 \times 10^{19} \text{ cm}^{-2}$. However, when smoothed to a WHAM resolution of 1 degree, the H I column density at this point is $\sim 1.0 \times 10^{19} \text{ cm}^{-2}$. Finally, towards (A7) there is a weak detection of ~ 0.25 R at $\sim +59$ km s^{-1} . At the WHAM resolution, the H I column density at this point is $\sim 0.4 \times 10^{19} \text{ cm}^{-2}$. We note that this position is close to the Arecibo-measured H I position ‘AR’ (Fig. 10), which has a similar velocity centroid of $v_{\text{LSR}} = +61$ km s^{-1} .




Figure 15. (a) Lovell-telescope IV H I column density map taken from Kennedy et al. (1998) in the vicinity of M15 with WHAM-observed positions marked (A0)–(A7). Contour levels (full line) are at $N_{HI} = (1, 2, 3) \times 10^{19} \text{ cm}^{-2}$. The dashed lines show the H α contours from the WHAM H α observations, integrated from $\sim +50$ – 90 km s^{-1} with levels at $0.1 \times (1, 2, 3, 4, 5) \text{ R}$. (b) H α brightness integrated from $\sim +50$ – 90 km s^{-1} for the whole field observed with the Lovell telescope in H I. Contour levels are at $0.1 \times (1, 2, 3, 4, 5) \text{ R}$. (c) WHAM H α spectra at the positions (A0)–(A7). Velocities are in the LSR. Where H α is observed (positions A0, A2, A5, A7), a three-component Gaussian fit is indicated by a dotted line with the (data–model) residual shown by the dashed line.

3.4.2 WHAM pointings towards HD 203664 and position ‘D’

The nearest WHAM H α pointings in the vicinity of the halo star HD 203664 are some 0.38 and 0.66 degrees from the stellar position and are depicted in Fig. 17. Although there is a slight excess of H α with velocities exceeding $\sim +50 \text{ km s}^{-1}$, this is very close to the noise and at lower velocities than the Ca II K absorption seen by Ryans et al. (1996) which had velocities of $+75$ – 80 km s^{-1} . Fig. 18 shows the WHAM pointings superimposed on the Lovell telescope multibeam H I column density map for the IVC towards IVC position ‘D’ (c.f. Fig. 10). Again, although there is some H α excess towards position (d3), this is very weak (brightness 0.07 R), and is also at a low-velocity of $\sim +50 \text{ km s}^{-1}$; this compares with the IVC H I velocity of $\sim +68 \text{ km s}^{-1}$ at this point. Even though the WHAM pointing centre is just within the $N_{HI} = 1 \times 10^{19} \text{ cm}^{-2}$ contour, the relatively small size of the

‘D’ component results in the H I column density at this point at 1 degree resolution being $\sim 0.5 \times 10^{19} \text{ cm}^{-2}$.

3.5 IRAS ISSA results

In this section the IRAS 60 and 100 micron ISSA images are compared with previous Arecibo H I observations of Smoker et al. (2001a). These latter data are of resolution 3 arcmin and contain both low and intermediate velocity gas. Fig. 19 shows the 100 micron image overlaid on the total (LV plus IV) H I column density, with Fig. 20 displaying the corresponding 60 micron data and the IVC H I column density alone. There appears to be a relatively good correlation between the IRAS and H I results, both for the low and intermediate velocity gas; although with such a small field such agreement could have occurred by chance. The emission is not likely to be from the M15 globular cluster itself, as glob-

Table 2. Wisconsin H α Mapper results. The values for the velocities and fluxes are the results of three-component Gaussian fitting to the low and intermediate velocity gas. H α brightnesses have not been corrected for Galactic absorption, hence assuming that the IVC lies above the Galactic plane, the tabulated values are likely to be low by ~ 25 per cent. Lovell telescope H I multibeam results, smoothed to a WHAM-sized 1 degree beam, provide H I velocity information towards position (A5).

Position on Fig. 15	(A7)	(A5)	(A2)	(A0)
RA (J2000)	$21^h 26^m 53.3^s$	$21^h 29^m 24.9^s$	$21^h 32^m 00.4^s$	$21^h 34^m 37.0^s$
Dec (J2000)	$+11^\circ 28' 58''$	$+12^\circ 14' 30''$	$+13^\circ 00' 02''$	$+13^\circ 44' 43''$
l (degrees)	63.88	64.98	66.09	67.19
b (degrees)	-27.16	-27.16	-27.16	-27.16
H α LV V_c (km s $^{-1}$)	$-32.0 \pm 16.6, +2.7 \pm 2.6$	$-53.9 \pm 4.6, -4.4 \pm 0.7$	$-43.8 \pm 3.7, -5.3 \pm 0.6$	$-32.4 \pm 9.5, -0.7 \pm 1.5$
H α LV FWHM (km s $^{-1}$)	$44.6 \pm 22.4, 35.1 \pm 4.8$	$26.7 \pm 10.4, 47.8 \pm 2.6$	$18.8 \pm 9.4, 35.2 \pm 2.0$	$30.3 \pm 14.1, 29.5 \pm 2.9$
H α LV Flux (R)	$0.27 \pm 0.18, 1.02 \pm 0.20$	$0.14 \pm 0.06, 2.10 \pm 0.08$	$0.11 \pm 0.04, 1.50 \pm 0.06$	$0.20 \pm 0.13, 1.26 \pm 0.14$
H α LV Peak (mR (km s $^{-1}$) $^{-1}$)	$4.8 \pm 1, 23 \pm 1$	$4.2 \pm 1, 35 \pm 1$	$4.7 \pm 0.1, 34 \pm 1$	$5.2 \pm 1, 34 \pm 1$
H α IV V_c (km s $^{-1}$)	$+59.3 \pm 1.1$	$+64.3 \pm 0.4$	$+59.4 \pm 0.8$	$+71.4 \pm 0.5$
H α IV FWHM (km s $^{-1}$)	18.8 ± 3.4	31.5 ± 1.4	38.1 ± 2.4	27.5 ± 1.5
H α IV Flux (R)	0.25 ± 0.02	1.30 ± 0.04	0.95 ± 0.04	0.91 ± 0.03
H α IV Peak (mR (km s $^{-1}$) $^{-1}$)	11 ± 1	33 ± 1	20 ± 1	26 ± 1
H I IV V_c (km s $^{-1}$)	—	65.7 ± 0.6	—	—
H I IV FWHM (km s $^{-1}$)	—	22.0 ± 2.0	—	—




Figure 16. Solid line: WHAM H α spectrum towards $l=64.98^\circ$, $b=-27.16^\circ$ (A5). The dotted line shows the results of a three-component Gaussian fit. The dashed line is a (data-three component Gaussian fit) residual.

ular clusters are undetected at the 60 and 100 micron IRAS sensitivity limits (Origlia, Ferraro & Pecci 1996). Plots comparing the low, total and intermediate H I column density with the IRAS flux density are displayed in Fig. 21. Assuming that the observed correlations are in fact real, the current data indicate that this IVC contains dust, as has also been observed in other such objects (Malawi 1989; Weiß et al. 1999). The fact that the IVC is tentatively detected at 60 microns indicates that the dust may be warm, perhaps heated by collisions.

The (tentative) IRAS detection contrasts with the situation in HVCs, which at least at the IRAS sensitivity do not appear to emit at these wavelengths (Wakker & Boulanger 1986). Such a difference indicates either differences in dust




Figure 17. Nearest two WHAM H α spectra towards HD 203664 with their angular distances from the star indicated.

content, dust temperature or environment (such as distance from the heating field of the Galactic plane). H I absorption line spectroscopy towards a number of HVCs by Akeson & Blitz (1999) indicates that the fraction of cold gas is low in these objects. If so, then this would point to the Galactic-halo HVCs containing less dust than the IVCs and not differences in temperature.

4 DISCUSSION

4.1 The H α properties of the M 15 IVC

4.1.1 Estimates of the emission measure and H II column density

The detection of H α towards the main M 15 IVC (position A5) with brightness of ~ 1.3 R compares to lower values of




Figure 18. Top panel: Lovell telescope multibeam IVC H I surface density map of feature (D) ($+50 < V_{LSR} < +80 \text{ km s}^{-1}$), with contour levels at $N_{HI} = (1, 2, 3) \times 10^{19} \text{ cm}^{-2}$ with WHAM pointings (d1)–(d5) marked. Bottom panel: WHAM H α spectrum towards (d3). The dotted line is a three-component Gaussian fit with the dashed line showing the data-fit residual spectrum.

between 0.06 to 0.20 R observed in a number of High Velocity Clouds (Tuftes et al. 1998). Similarly, many of the HVCs observed by Weiner, Vogel & Williams (1999) are very faint ($< 0.03 \text{ R}$), implying that, if they are photoionised by hard photons from the Galaxy, they are 20–60 kpc distant. The relatively high H α emission from the M15 IVC in turn argues for a closer distance, assuming that shock ionisation is not the dominant factor. This is a big assumption, as detections of H α in the Magellanic Stream (distance 55 kpc) of are 0.20–0.37 Rayleigh (Bland-Hawthorn & Maloney 1999) are similar to the range of ~ 0.1 –0.5 Rayleigh detected towards the IVC Complex K (Haffner, Reynolds & Tuftes 2001a) which has a distance bracket of 0.3–7.7 kpc.

Assuming that extinction is negligible and that the cloud is optically thin, the observed H α brightness of the IVC can be used to estimate both the column density of




Figure 19. Arecibo H I and IRAS ISSA data. Greyscale: IRAS 100 micron flux density. Contour: Total (LV plus IV) H I column density at $N_{HI} = (56, 60, 64, 68, 72) \times 10^{19} \text{ cm}^{-2}$.




Figure 20. Arecibo H I and IRAS ISSA data. Greyscale: IRAS 60 micron flux density. Contour: IVC H I column density at $N_{HI} = (2, 4, 6, 8) \times 10^{19} \text{ cm}^{-2}$. Note how the peak H I contour corresponds with the IRAS peak.

ionised hydrogen and the electron density. For an optically thin gas, the H α surface brightness in Rayleigh, I_{α} , is given by;

$$I_{\alpha} = 0.36 \times \text{EM} \times \sqrt{T} \times (1 - 0.37 \times \ln(T)), \quad (1)$$

where T is the gas temperature in units of 10^4 K and EM is




Figure 21. Arecibo H I and IRAS ISSA data. The IRAS fluxes have an arbitrary zero-point. (a) IRAS 100 micron flux density plotted against low velocity gas H I column density. (b) IRAS 100 micron flux density plotted against total H I column density. (c) IRAS 60 micron flux density plotted against IVC H I column density alone.

the emission measure ($\int n_e^2 dl$) in units of pc cm^{-6} (Simonetti, Topasna & Dennison 1996). Here, n_e is the electron density and integration over dl provides the thickness (L) of the ionised region. Assuming the temperature of the H II is 8,000 K, typical of the warm interstellar medium (Reynolds 1985), the resulting emission measure towards position (A5) is 3.7 pc cm^{-6} .

Assuming a distance to the main IVC condensation of 1 kpc, the cloud size (d) at a column density limit of $N_{\text{HI}} = 10^{19} \text{ cm}^{-2}$ is $\sim 0.6^\circ \times 0.8^\circ$, corresponding to linear dimensions of $\sim 10 \times 14 \text{ pc}$. If the cloud is approximately spherically symmetric, this results in an estimated electron density (assuming a size of 12 pc) of $\sqrt{\text{EM}/d} \sim 0.6 f^{-1/2} \text{ cm}^{-3}$ and a column density of ionised hydrogen ($= L \times n_e$) of $\sim 2 \times 10^{19} f^{1/2} \text{ cm}^{-2}$, where f is the filling fraction of H II in the IVC. Smoothed to a resolution of 1 degree (corresponding to the size of the WHAM beam), the H I column density at this point derived from the data of Kennedy et al. (1998) is $\sim 1.4 \times 10^{19} \text{ cm}^{-2}$, hence the fractional ratio of H II to H I is of order $1.4 \times f^{1/2}$, indicating that there is a substantial amount of ionised gas at intermediate velocities present in this sightline.

For positions (A2) and (A7), and using the same cloud size as above, results in estimated H II column densities of $1.5 \times 10^{19} \text{ cm}^{-2}$ and $0.8 \times 10^{19} \text{ cm}^{-2}$, corresponding to frac-

tional H II to H I values of order $1.5 \times f^{1/2}$ and $2.0 \times f^{1/2}$ respectively. These fractional H II to H I ratios towards the current IVC are somewhat higher than derived by Tufte et al. (1998) for a sample of HVCs which were calculated to be $< 0.06 \times f^{1/2}$. However, recent work by Bluhm et al. (2001), in sightlines towards the Large Magellanic Cloud, used the relative underabundance of neutral oxygen to infer an ionisation level in both an IVC and HVC of ~ 90 per cent. It would be useful to observe this cloud using the same methods as described in the current paper and compare the results.

Towards (d3) there is no detection in H α with velocities corresponding to the H I value of $\sim +68 \text{ km s}^{-1}$. Assuming a cloud size of 8 pc and upper limit to the H α brightness of 0.2 R, gives an estimated upper limit to the IVC H II column density and fractional H II to H I values of $\sim 4 \times 10^{18} \text{ cm}^{-2}$ and $0.7 \times f^{1/2}$. Finally, we consider the halo star HD 203664, towards which the limiting column density of neutral hydrogen is $\sim 10^{18} \text{ cm}^{-2}$. If we take the upper limit to the H α brightness of 0.2 R, and combine this with a cloud size L (in pc, where the Arecibo beam is $\sim 1 \text{ pc}$ FWHM at 1 kpc distance), a fractional H II to H I ratio of $\sim 1.4 \times L \times f^{1/2}$ can be set by the current observations.

4.1.2 Estimates of the ionising radiation field and electron density

Following Tufte et al. (1998), if photoionisation is the dominant cause of H α emission, then the incident Lyman continuum flux F_{LC} can be estimated thus, assuming case B recombination;

$$F_{\text{LC}} = 2.1 \times 10^5 \frac{I_\alpha}{0.1\text{R}} \text{ photons cm}^{-2} \text{ s}^{-1}, \quad (2)$$

-where I_α is the H α intensity in Rayleigh. For the M 15 IVC, equation 2 implies an incident flux (F_{LC}) of $2.7 \times 10^6 \text{ photons cm}^{-2} \text{ s}^{-1}$. Hence if photoionisation is the main cause of H α production, the derived Lyman-alpha continuum flux towards the main M 15 IVC condensation is a factor 6–22 times higher than the implied incident flux towards the A, C and M HVCs observed by Tufte et al. (1998), and more than twice that observed towards the Complex K IVC by Haffner et al. (2001a). We recall that Complex A lies between 4–10 kpc, with Complex C being some 5–25 kpc distant and the M 15 IVC being closer than 3 kpc. In the future, comparison of derived Lyman-alpha continuum fluxes for a larger sample of IVC and HVC sightlines with known distances may provide information on the relative contributions of the Galactic and extragalactic ionising field.

An alternative possibility is that the H α is produced by shock ionisation, caused by interaction of the IVC with LV gas. This is a real possibility given the orientation of the IVC and the fact that its z -distance of less than 1 kpc puts it in the lower Galactic halo. Towards the nearby halo star HD 203664 in which IV absorption is seen, Sembach (1995) postulated that the dust grains in the IVC have been processed by such shocks which also currently produce the highly ionised species. For an ambient temperature of $< 3 \times 10^5 \text{ K}$, a cloud of velocity 50 km s^{-1} will be supersonic and hence shocks may be formed given the right conditions. Using the models of Raymond (1979), which are applicable for shocks with speed $50 < V_s < 140 \text{ km s}^{-1}$, the face-on

H α surface brightness ($I_{\alpha\text{perp}}$) can be related to the number density of the pre-shocked gas (n_0), thus;

$$(I_{\alpha\text{perp}}) \sim 6.5 \times n_0 (V_s/100)^{1.7} R \quad (3)$$

Given that $I_{\alpha\text{perp}} = 1.3 R$, and assuming a shock speed of 50 km s^{-1} , leads to an upper limit to n_0 of 0.6 cm^{-3} . The value is an upper limit (for this shock speed velocity) as a non-perpendicular sightline will increase the observed I_{α} (Tufte et al. 1998). Of course, given the fact that the transverse component of the velocity of the M 15 IVC is unknown, this value is very uncertain. Discriminating between shock and photoionisation is difficult, although further progress may be possible via measurements of appropriate emission line ratios (Tufte et al. 1998).

4.2 Ca II number density towards the M 15 IVC and HD 203664

Before discussing the Ca II K results, we note that calcium is depleted onto dust and is not the dominant ionisation species, hence the absolute metallicity of the M 15 IVC is uncertain and awaits high resolution UV observations. As emphasized by Ryans et al. (1997), differences in resolution between the optical and radio data, combined with the Ca II K results only placing limits on the ion abundances, makes it important not to over-interpret the observed $N(\text{Ca II})/N(\text{H I})$ ratios.

With that caveat, *and assuming that the current observations do not miss any narrow-velocity components*, the average IVC Ca II number density and ratio of the IVC compared to H I towards the centre of M 15 are $\log_{10}(N(\text{Ca II}) \text{ cm}^{-2}) = 12.0$ and $N(\text{Ca II})/N(\text{H I}) = 2.5 \times 10^{-8}$. The H I column density towards the M 15 centre is $4 \times 10^{19} \text{ cm}^{-2}$ and was obtained using the combined WSRT plus Arecibo map of resolution $111'' \times 56''$. For the nearby sightline towards the halo star HD 203664, we use the results of Ryans et al. (1996) for the total IVC Ca II K column density of $\sim 1 \times 10^{12} \text{ cm}^{-2}$, combined with the upper limit to the H I column density at a resolution of 3 arcmin towards HD 203664 of $\sim 1 \times 10^{18} \text{ cm}^{-2}$ (Smoker et al. 2001a), giving a much higher value of $N(\text{Ca II})/N(\text{H I}) > 1.0 \times 10^{-6}$.

The current results compare with literature values of $N(\text{Ca II})/N(\text{H I}) \sim 2 \times 10^{-9}$ (Hobbs 1974, 1976) for low velocity diffuse clouds and of $N(\text{Ca II})/N(\text{H I})$ in the range $3\text{--}300 \times 10^{-9} \text{ cm}^{-2}$ for the high latitude clouds studied by Albert et al. (1993). The $N(\text{Ca II})/N(\text{H I})$ ratios in IVCs and HVCs are thought to be higher than in low velocity gas due to the former having less dust onto which calcium is depleted. Thus the observed $N(\text{Ca II})/N(\text{H I})$ ratio of 2.5×10^{-8} (or ~ 0.01 of the total solar calcium abundance) in the line-of-sight towards M 15 is typical of other high latitude clouds and also of other HVCs and IVCs (e.g. Wakker et al. 1996a; Ryans et al. 1997).

The lower limit of $N(\text{Ca II})/N(\text{H I}) > 10^{-6}$ towards HD 203664 is, however, on the high side for IVCs/HVCs, being ~ 0.5 of the total solar calcium abundance. There are several possible reasons for this. Firstly, the (currently undetected) H I towards HD 203664 could be in a clump of gas smaller than the Arecibo beamsize of 3 arcmin; if this were the case then the H I column density limit used would be too low and the derived $N(\text{Ca II})/N(\text{H I})$ ratio too high.

Additionally, HST UV observations towards HD 203664 indicate that the H I towards this object is at least partially ionised (Sembach, private communication), either by shock ionisation or photoionisation. If photoionisation, aside from the normal ionising source being Galactic OB-type stars or the extragalactic UV field, HD 203664 itself (spectral type B0.5) could be a possible ionising source. The fact that its LSR velocity is $+110 \text{ km s}^{-1}$ (Little et al. 1994) compared with the IVC at $+70 \text{ km s}^{-1}$ is inconclusive in determining the relative distance of the line of sight IVC towards HD 203664 with the star itself. Finally, there remains the possibility that the hydrogen is in molecular form. However, given the low H I column density towards the HD 203664 sightline, this appears unlikely.

Alternatively, it could be that the derived value of $N(\text{Ca II})/N(\text{H I}) > 10^{-6}$ towards HD 203664 is correct. This would tally with the IUE results of Sembach (1995), which found that the majority of the elements in the IV gas, when referenced to sulphur, were within a factor 5 of their solar values and strongly point to a Galactic origin for this part of the IVC. The fact that our derived value for $N(\text{Ca II})/N(\text{H I})$ towards the M 15 IVC of 0.01 solar is much lower than towards HD 203664 is likely to be caused by different ionisation fractions and dust contents, and/or differing formation mechanisms. Clearly the latter is speculative and requires follow-up high resolution UV observations towards the M 15 IVC to determine the abundances of elements such as sulphur and zinc that are not depleted onto dust.

4.3 Velocity widths and temperatures towards the M 15 IVC and HD 203664

Towards the main M 15 IVC condensation, (feature ‘A’ on Fig. 10) values of the H I FWHM velocity width at resolutions of $\sim 2 \times 1$ arcmin range from $5\text{--}14 \text{ km s}^{-1}$, corresponding to maximum kinetic temperatures in the range $\sim 500\text{--}4000 \text{ K}$. Mid-way between the M 15 IVC and HD 203664, the FWHM equals 12 km s^{-1} at a resolution of 3 arcmin, which corresponds to $T_k \sim 3000 \text{ K}$ (feature ‘AR’ on Fig. 10). The current observations have additionally observed feature ‘D’ at a resolution 12 arcmin, with FWHM velocity width also of 12 km s^{-1} , indicating gas of similar temperature. We note that each of these temperatures will be upper limits due to beamsmeaking and turbulent velocity components. Finally, towards HD 203664, Ryans et al. (1996) found cloudlets with FWHM of 2.8 and 3.2 km s^{-1} in Ca II K, corresponding to upper limits for the kinetic temperatures of $\sim 8000\text{--}10,000 \text{ K}$. Towards the same star, Sembach (1995) used the relative abundances of low-ionisation species to derive a temperature for the HD 203664 IVC of $5300\text{--}6100 \text{ K}$.

The higher IVC gas temperatures towards HD 203664 than towards the M 15 IVC, feature ‘D’ or the intermediate position ‘AR’ could be interpreted as being caused by the former cloud being nearer to the ionising field of the Galactic plane than the other two IVCs (Lehner 2000). It seems more likely that the lower temperatures seen towards parts of ‘A’ and ‘AR’ are simply caused by shielding of parts of these gas clouds; shielding that is not possible towards the HD 203664 IVC because of the lower gas density there. A two-phase core-envelope structure for halo HVCs *has* often been proposed within the Galactic corona for $1 < z < 5 \text{ kpc}$

(e.g. Wolfire et al. 1995), where the two components of 100 and 10,000 K are entrained by pressure from the hot Galactic corona. We note that the high-end H I temperatures observed towards the M 15 IVC indeed occur towards its outer parts where the H I column density is low and the FWHM velocity widths are uncertain.

4.4 Comparison of H I properties with previous Galactic halo IVCs and HVCs

In this section we compare the high-resolution H I properties of the M 15 IVC with other IVCs and HVCs known to lie within the Galactic halo and observed at comparable resolution. We note that it is likely that there are many different types of HVC, with recent work, for example, indicating that a number of the compact HVCs lie at distances of several tens of kpc (Braun & Burton 2000). A literature search found the following objects with known distances and observed in H I at high resolution: the M 13 IVC (Shaw et al. 1996), the 4-Lac HVC 100-7+100 (Stoppelenburg et al. 1998), the Low-Latitude Intermediate Velocity Arch (Wakker et al. 1996b), IVC 135+54-45 (Weiß et al. 1999), HVC 132+23-211 (within Complex A; Schwarz et al. 1976) and the M 92 HVC (within Complex C; Smoker et al. 2001b).

Table 3 compares the properties of these IVC and HVC H I gas clouds known to exist in the Galactic halo. Inspecting the table, there are no clear differences in the H I properties (column density, peak temperature) of the two types of objects, which show a large scatter both within IVCs and HVCs. Similarly, the velocity widths of all of the objects, barring the peculiar HVC100-7 and the Mirabel cloud, all show minimum values of FWHM $\sim 5 \text{ km s}^{-1}$ and indicating that gas of temperature less than $\sim 500 \text{ K}$ is common in these objects. This is in contrast to the situation observed at lower resolution for some northern clouds, where IVCs tend to have smaller velocity widths than their HVC counterparts (Davies, Buhl & Jafolla 1976).

If some IVCs and HVCs are formed from infalling gas, sweeping up high- z H I as they fall towards the plane, or if they are objects within the plane, or if they are formed within a Galactic fountain, it seems plausible that they could be preferentially aligned with the Galactic plane. Of course, it must be taken into consideration that at high resolution, only small parts of cloud complexes are studied, and by chance, some of these components will be aligned with the plane. In any case, of the four objects in the sample with a well-defined axis and at high Galactic latitude, three have their major axis near-parallel with the plane. Although there exist a number of such objects observed at lower resolution with this orientation, to our knowledge no systematic survey has been performed determining the orientation parameters of HVCs. If performed, this could act as a further discriminator between HVCs known to exist in the Galactic halo, and the sample of HVCs postulated to lie at extragalactic distances.

Summarising, at present there are too few high-resolution observations of Galactic halo IVCs and HVCs to determine differences in H I properties and any relationship between the two types of object. However, as previously noted, the IRAS and H α properties *do* appear to differ, al-

though the number of objects studied in all three wavebands remains small.

5 SUMMARY AND CONCLUSIONS

The current H I WSRT synthesis observations have shown that on scales down to ~ 1 arcmin, the M 15 IVC shows substructure, with variations in the column density of a factor of ~ 4 on scales of ~ 5 arcmin being observed, corresponding to scales of $\sim 1.5 D$ pc, where D is the the IVC distance in kpc. Of course, this is not an unexpected finding, but once again demonstrates that great care must be taken in interpreting quantities such as cloud metallicities which are derived from a combination of low-resolution radio plus optical data. The Lovell telescope H I observations towards this cloud demonstrated how relatively large areas of sky can be mapped with the multibeam system in a short period of time in the search for IVCs and HVCs. These data showed that the M 15 IVC has components spread out over several square degrees, with component ‘D’ being mapped for the first time at medium resolution (12 arcmin) and having a similar column density to the IV gas centred upon M 15 itself. Both the H I emission-line and Ca II absorption-line data showed *tentative* evidence for velocity substructure, perhaps indicative of cloudlets. The Ca II/H I value of $\sim 2.5 \times 10^{-8}$ towards the main M 15 condensation is similar to that previously observed in other IVCs and HVCs. Towards HD 203664, the *observed* lower limit of 10^{-6} is somewhat higher, although this may be caused by factors such as the H I beam being unfilled or partial ionisation of the gas on this sightline. The H I properties of the M 15 IVC are indistinguishable from HVCs, although with the lack of distance information towards most HVCs, comparisons are difficult.

The tentative detection of infrared emission from the M 15 IVC, as in other IVCs, does distinguish it from HVCs, and either points to the M 15 IVC containing more dust, and/or being closer to the heating field of the Galactic plane than HVCs, which as a class of objects are not detected in the IRAS wavebands. Similarly, the relatively strong H α emission (exceeding 1 Rayleigh) towards parts of the M 15 IVC, *if* caused by photoionization, may place it closer to the Galaxy than HVCs. Again, however, this finding is uncertain due to the problem in distinguishing photoionisation from shock ionisation, uncertainties in dust content, and differences in H I volume densities in different objects studied thus far.

Future work towards this cloud should include higher-signal-to-noise observations in the Ca II line in order to determine if the cloud velocity substructure tentatively found in the current observations is in fact real, and whether the Ca II/H I ratio determined by the current observations is lower than towards the HD 203664 sightline. This should be combined with $^{12}\text{CO}(1-0)$ sub-mm observations in order to determine if molecular material exists towards the peaks in H I column density and out of which stars may form. The determination of the falloff in H I column density, of the cloud to low column density limits would also indicate the ionisation properties of the object and whether or not there is any interaction between the M 15 IVC gas and low velocity material. Finally, UV observations towards M 15 globular cluster stars, although difficult, would provide important in-

Table 3. Previous high-resolution H I observations of IVCs and HVCs known to exist in the halo, compared with the current results. T_p and N_{HI} are the peak temperatures in K and peak H I column densities in units of 10^{19} cm^{-2} respectively. $\Delta V1$ and $\Delta V2$ are the broad and narrow velocity widths in km s^{-1} . The field “Parallel to the Gal. plane” only refers to the object observed at high-resolution and not to the whole complex in the case of the large HVCs. References; RKSD97, Ryans et al. (1997); SSH76, Schwarz et al. (1976); SBK96, Shaw et al. (1996); SSW98, Stoppelenburg et al. (1998); SKD01, Smoker et al. (2001b); WHS96, Wakker et al. (1996b); WPS99, van Woerden et al. (1999); WWHM99 Weiß et al. (1999). GSR velocities are calculated via $V_{GSR} = V_{LSR} + 250 \times \sin(l) \times \cos(b) \text{ km s}^{-1}$.

Cloud Ref.	l° b°	D (kpc)	V_{LSR} V_{GSR}	$\Delta V1$ $\Delta V2$	T_p N_{HI}	IRAS?	Ca II/H I	Parallel to Gal. plane?	Res. '
HVC 132+23–211 (SSH76, WPS99)	123 +23	4–10	–211 –329	5–15	25 26	N	–	Y	~ 2
M13SE IVC (SBK96)	59 +41	<7	–73 +89	27–35 5	0.7 4.7	N	$>1 \times 10^{-7}$	N	$\sim 3 \times 2$
M15 IVC (This paper)	65 –27	<3	+70 +272	5–15 –	8 15	Y?	2.5×10^{-8}	Y	$\sim 2 \times 1$
M92N HVC (SKD01)	68 +34	5–25	–101 +91	4–7 –	3.4 6	N	–	Y	$\sim 6 \times 6$
HVC100–7+100 (SSW98)	100 –7	<1.2	+106 +350	13 –	0.5 0.16	N	–	N	$\sim 2 \times 2$
IVC 135+54–45 (WHM99)	135 +54	0.29–0.39	–45 +59	4–5 –	>7 30?	Y	–	–	$\sim 1 \times 1$
PG0859+593 LLIVC (RKSD97, WHS96b)	+156.9 +39.7	1.7–4.0	–51 +24	+19	2.0 5.3	?	–	–	$\sim 2 \times 2$
PG0906+597 LLIVC (RKSD97, WHS96b)	+156.2 +40.6	1.7–4.0	–48, –53 +29, +24	28, 6	1.6, 3.9 11.0, –	?	–	–	$\sim 2 \times 2$

formation on the absolute metallicity of the gas towards this object for comparison with the HD 203664 sightline.

ACKNOWLEDGEMENTS

We would like to thank the Netherlands Foundation for Research in Astronomy (NFRA) which is a national facility supported by the Netherlands Organization for Scientific Research. In particular, we are very grateful for the help that Robert Braun provided with the reduction of the WSRT data. JVS acknowledges NFRA for use of data reduction facilities and hospitality and to the staff of Jodrell Bank observatory for help with the Lovell telescope multibeam observations. We would particularly like to thank Chris Jordan, Rob Lang, Peter Boyce and Robert Minchin for help with the LT observations. JVS would also like to thank the staff of the William Herschel Telescope, which is part of the Isaac Newton Group of telescopes, La Palma. The Digitized Sky Surveys were produced at the Space Telescope Science Institute under U.S. Government grant NAG W-2166. The National Geographic Society - Palomar Observatory Sky Atlas (POSS-I) was made by the California Institute of Technology with grants from the National Geographic Society. IPAC is operated by the Jet Propulsion Laboratory (JPL) and California Institute of Technology (Caltech) for NASA. IPAC is funded by NASA as part of the IRAS extended mission program under contract to JPL/Caltech. The WHAM project is funded primarily through grants from the National Science Foundation with additional support provided by the University of Wisconsin. JVS also thanks Andrew George and PPARC for financial support and to the referee, Dr V. Kilborn for useful comments.

REFERENCES

- Akeson R. L., Blitz L., 1999, *ApJ*, 523, 163
 Albert C. E., Blades J. C., Morton D. C., Lockman F. J., Proulx M., Ferrarese L., 1993, *ApJS*, 88, 81
 Allen C. W., 1973, *Astrophysical Quantities*, 3rd edition, University of London, Athlone Press
 Barnes D. G., Staveley-Smith L., de Blok W. J. G., Oosterloo T., Stewart I. M., Wright A. E., et al., 2001, *MNRAS*, 322, 486
 Bland-Hawthorn J., Maloney P. R., 1999, *ApJ*, 510, 33
 Blitz L., Spergel D. N., Teuben P. J., Hartmann D., Burton W. Butler, 1999, *ApJ*, 514, 818
 Bluhm H., de Boer K. S., Marggraf O., Richter P., 2001, *A&A*, 367, 299
 Bowen D. V., 1991, *MNRAS*, 251, 649
 Braun R., Burton W. B., 2000, *A&A*, 354, 853
 Brüns C., Kerp J., Kalberla P. M. W., Mebold U., 2000, *A&A*, 357, 120
 Cardelli J. A., Clayton G. C., Mathis J. S., 1989, *ApJ*, 345, 245
 Christodoulou D. M., Tohline J. E., Keenan F. P., 1997, *ApJ*, 486, 810
 Davies R. D., Buhl D., Jafolla J., 1976, *A&AS*, 23, 181
 Durrell P. R., Harris W. E., 1993, *AJ*, 105, 1420
 Dyson J. E., Hartquist T. W., 1983, *MNRAS*, 203, 1233
 Faison M. D., Goss W. M., Diamond P. J., Taylor G. B., 1998, *AJ*, 116, 2916
 Haffner L. M., 1999, PhD Thesis, University of Wisconsin–Madison, U.S.A.
 Haffner L. M., Reynolds R. J., Tufte S. L., 2001a, *ApJL*, 556, 33
 Haffner L. M., Reynolds R. J., Tufte S. L., et al., 2001b, in preparation
 Harris W. E., 1996, *AJ*, 112, 1487
 Hartmann D., Burton W. B., 1997, *Atlas of galactic neutral hydrogen*, Cambridge University Press
 Heiles C., 1997, *ApJ* 481, 193
 Hobbs L. M., 1974, *ApJ*, 191, 381
 Hobbs L. M., 1976, *ApJ*, 203, 143
 Howarth I. D., Murray J., Mills D., Berry D. S., 1996, *STAR-LINK, User Note SUN 50*, Rutherford Appleton Labora-

- tory/CCLRC
- Ivezic Z., Christodoulou D. M., 1997, *ApJ*, 486, 818
- Jaschek C., Jaschek M., 1987, *The classification of stars*, Cambridge University Press
- Kennedy D. C., Bates B., Keenan F. P., Kemp S. N., Ryans R. S. I., Davies R. D., Sembach K. R., 1998, *MNRAS*, 297, 849
- Kuntz K. D., Danly L., 1996, *ApJ*, 457, 703
- Lauroesch J. T., Meyer D. M., Blades J. C., 2000, *ApJ*, 543, 43
- Lehner N., Rolleston W. R. J., Ryans R. S. I., Keenan F. P., Bates B., Pollacco D. L., Sembach K. R., 1999, *A&AS*, 134, 257
- Lehner N., 2000, PhD Thesis, The Queen's University of Belfast
- Little J. E., Dufton P. L., Keenan F. P., Conlon E. S., Davies R. D., 1994, *ApJ*, 427, 267
- Malawi A., 1989, PhD Thesis, The University of Manchester
- Meyer D. M., Lauroesch J. T., 1999, *ApJ*, 520, 103
- Morton D. C., 1991, *ApJS*, 77, 119
- Origlia L., Ferraro F. R., Pecci F. F., 1996, *MNRAS*, 280, 572
- Raymond J. C., 1979, *ApJS*, 39, 1
- Reynolds R. J., 1985, *ApJ*, 294, 256
- Ryans R. S. I., Sembach K. R., Keenan F. P., 1996, *A&A*, 314, 609
- Ryans R. S. I., Keenan F. P., Sembach K. R., Davies R. D., 1997, *MNRAS*, 289, 83
- Sault R. J., Teuben P. J., Wright M. C. H., 1995, *Astronomical Data Analysis Software and Systems IV*, ed. Shaw R., Payne H. E., Hayes J. J. E., ASP Conf. Ser. 77, 433
- Schwarz U. J., Sullivan W. T. III., Hulsbosch A. N. M., 1976, *A&A*, 52, 133
- Sembach K. R., 1995, *ApJ*, 445, 314
- Shaw C. R., Bates B., Kemp S. N., Keenan F. P., Davies R. D., Roger R. S., 1996, *ApJ*, 473, 849
- Shortridge K., Meyerdierks H., Currie M., Clayton M., Lockley J., Charles A., et al., 1999, *STARLINK, User Note SUN 86*, Rutherford Appleton Laboratory/CCLRC
- Simonetti J. H., Topasna G. A., Dennison B., 1996, *AAS*, 188, 12.01
- Smoker J. V., Lehner N., Keenan F. P., Totten E. J., Murphy E., Sembach K. R., Davies R. D., Bates B., 2001a, *MNRAS*, 322, 13
- Smoker J. V., Roger R. S., Keenan F. P., Davies R. D., Lang R. H., 2001b, *A&A*, 380, 673
- Stoppelenburg P. S., Schwarz U. J., van Woerden H., 1998, *A&A*, 338, 200
- Tan J. C., 2000, *ApJ*, 536, 173
- Tufte S. L., Reynolds R. J., Haffner L. M., 1998, *ApJ*, 504, 773
- van Woerden H., Peletier R. F., Schwarz U. J., Wakker B. P., Kalberla P. M. W., 1999, In *Stromlo Workshop on High-Velocity Clouds*, eds. Gibson B. K., Putman M. -E., ASP Conference Series Vol. 166, 1
- Wakker B. P., Boulanger F., 1986, *A&A*, 170, 84
- Wakker B. P., 1991, *A&A*, 250, 499
- Wakker B. P., Schwarz U. J., 1991, *A&A*, 250, 484
- Wakker B. P., van Woerden H., Schwarz U. J., Peletier R. F., Douglas N. G., 1996a, *A&A*, 306, 25
- Wakker B., Howk C., Schwarz U., van Woerden H., Beers T., Wilhelm R., Kalberla P., Danly L., 1996b, *ApJ*, 473, 834
- Wakker, B. P. 2001, *ApJS*, 136, 463
- Weiβ A., Heithausen A., Herbstmeier U., Mebold U., 1999, *A&A*, 344, 955
- Weiner B. J., Vogel S. N., Williams T. B., 1999, *BAAS*, 195, 9703 (abstr.)
- Wolfire M. G., McKee C. F., Hollenbach D., Tielens A. G. G. M., 1995, *ApJ*, 453, 673